



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINES  
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 5/19/2016  
LAST REVIEW DATE: 5/19/2022  
LAST CRITERIA REVISION DATE: 5/19/2022  
ARCHIVE DATE:

---

## CRESEMBA® (isavuconazonium sulfate) oral

---

Coverage for services, procedures, medical devices and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Pharmacy Coverage Guideline must be read in its entirety to determine coverage eligibility, if any.

This Pharmacy Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide BCBSAZ complete medical rationale when requesting any exceptions to these guidelines.

The section identified as "**Description**" defines or describes a service, procedure, medical device or drug and is in no way intended as a statement of medical necessity and/or coverage.

The section identified as "**Criteria**" defines criteria to determine whether a service, procedure, medical device or drug is considered medically necessary or experimental or investigational.

State or federal mandates, e.g., FEP program, may dictate that any drug, device or biological product approved by the U.S. Food and Drug Administration (FDA) may not be considered experimental or investigational and thus the drug, device or biological product may be assessed only on the basis of medical necessity.

Pharmacy Coverage Guidelines are subject to change as new information becomes available.

For purposes of this Pharmacy Coverage Guideline, the terms "experimental" and "investigational" are considered to be interchangeable.

**BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.**

---

This Pharmacy Coverage Guideline does not apply to FEP or other states' Blues Plans.

Information about medications that require precertification is available at [www.azblue.com/pharmacy](http://www.azblue.com/pharmacy).

Some large (100+) benefit plan groups may customize certain benefits, including adding or deleting precertification requirements.

All applicable benefit plan provisions apply, e.g., waiting periods, limitations, exclusions, waivers and benefit maximums.

Precertification for medication(s) or product(s) indicated in this guideline requires completion of the [request form](#) in its entirety with the chart notes as documentation. **All requested data must be provided.** Once completed the form must be signed by the prescribing provider and faxed back to BCBSAZ Pharmacy Management at (602) 864-3126 or emailed to [Pharmacyprecert@azblue.com](mailto:Pharmacyprecert@azblue.com). **Incomplete forms or forms without the chart notes will be returned.**



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINES  
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 5/19/2016  
LAST REVIEW DATE: 5/19/2022  
LAST CRITERIA REVISION DATE: 5/19/2022  
ARCHIVE DATE:

---

## CRESEMBA® (isavuconazonium sulfate) oral

---

### Criteria:

- **Criteria for initial therapy:** Cresemba (isavuconazonium sulfate) is considered *medically necessary* and will be approved when **ALL** of the following criteria are met:
1. Prescriber is a physician specializing in the patient's diagnosis or is in consultation with an Infectious Disease Specialist, Transplant Physician, or Oncologist
  2. Individual is 18 years of age or older
  3. **For a diagnosis of invasive aspergillosis**, individual has failure, contraindication or intolerance to **BOTH** of the following:
    - a. Vfend (voriconazole)
    - b. Posaconazole (Noxafil or generic) tablet
  4. **For a diagnosis of invasive mucormycosis**, individual has failure, contraindication or intolerance to **BOTH** of the following:
    - a. Lipid formulation of amphotericin B (either liposomal amphotericin B or amphotericin B lipid complex)
    - b. Posaconazole (Noxafil or generic) tablet
  5. **ALL** of the following baseline tests have been completed before initiation of treatment with continued monitoring as clinically appropriate:
    - a. Fungal culture report shows causative organism(s) are sensitive to isavuconazonium only
  6. There are **NO** FDA-label contraindications, such as:
    - a. Use with strong CYP3A4 inhibitors, such as ketoconazole or high-dose ritonavir (400 mg every 12 hours)
    - b. Use with strong CYP3A4 inducers, such as rifampin, carbamazepine, St. John's wort, or long acting barbiturates
    - c. Use in patients with familial short QT syndrome

**Initial approval duration:** 3 months

- **Criteria for continuation of coverage (renewal request):** Cresemba (isavuconazonium sulfate) is considered *medically necessary* and will be approved when **ALL** of the following criteria are met:
1. Individual continues to be seen by a physician specializing in the patient's diagnosis or is in consultation with an Infectious Disease Specialist, Transplant Physician, or Oncologist
  2. Individual's condition responded and has not worsened while on therapy
    - a. Response is defined as **TWO** of the following:
      - i. Clinical signs and symptoms of infection are resolving
      - ii. Radiographic abnormalities are improving



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINES  
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 5/19/2016  
LAST REVIEW DATE: 5/19/2022  
LAST CRITERIA REVISION DATE: 5/19/2022  
ARCHIVE DATE:

---

## CRESEMBA® (isavuconazonium sulfate) oral

---

- iii. There is no evidence of disease progression
- b. Worsening is defined as any of the following:
  - i. Signs and symptoms of the infection have not resolved
  - ii. Radiographic abnormalities continue to progress
- 3. The indication for use is one that requires a longer duration than the usual duration
- 4. Individual has been adherent with the medication
- 5. Individual has not developed any contraindications or other significant adverse drug effects that may exclude continued use
  - a. Contraindications as listed in the criteria for initial therapy section
  - b. Significant adverse effect such as:
    - i. Liver toxicity
    - ii. Exfoliative cutaneous reactions or Stevens Johnson syndrome
- 6. There are no significant interacting drugs

**Renewal duration:** 3 months

➤ Criteria for a request for non-FDA use or indication, treatment with dosing, frequency, or duration outside the FDA-approved dosing, frequency, and duration, refer to one of the following Pharmacy Coverage Guideline:

1. **Off-Label Use of Non-Cancer Medications**
2. **Off-Label Use of Cancer Medications**

---

### **Description:**

Cresemba (isavuconazonium) is an azole antifungal medication indicated for use in individuals 18 years of age or older for the treatment of invasive aspergillosis and invasive mucormycosis. Isavuconazonium is a pro-drug that is converted to its active component, isavuconazole.

Aspergillus is a life-threatening infection seen in immunocompromised patients. Invasive aspergillosis is an important cause of mortality and morbidity in patients with prolonged neutropenia, hematologic malignancies, and allogeneic hematopoietic stem cell transplantation (HSCT) recipients.

Mucormycosis is a very aggressive invasive fungal disease. Invasive mucormycosis is a rare, life-threatening mold infection, usually seen in immunocompromised patients, such as those with hematologic malignancies and in HSCT recipients.



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINES  
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 5/19/2016  
LAST REVIEW DATE: 5/19/2022  
LAST CRITERIA REVISION DATE: 5/19/2022  
ARCHIVE DATE:

---

## CRESEMBA® (isavuconazonium sulfate) oral

---

Isavuconazole has activity against most strains of the following microorganisms, both *in vitro* and in clinical infections: *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus niger*, and Mucorales such as *Rhizopus oryzae* and Mucormycetes species.

Studies suggest cross-resistance between isavuconazole and other azoles. The relevance of cross-resistance to clinical outcome has not been fully characterized. Individuals failing prior azole therapy may require alternative antifungal therapy.

Empiric therapy with an antifungal that has adequate coverage against the suspected causative organism is recommended until the diagnosis is confirmed. Definitive choice of systemic antifungal therapy is based on sensitivity data from fungal cultures, identification of causative organism, site of infection, immune status of the patient, and other patient characteristics. Once these results of the laboratory tests become available, antifungal therapy should be adjusted accordingly.

An Infectious Diseases Society of America (IDSA) summary guideline provides a comprehensive, evidence-based review of antifungal therapy for the management of aspergillosis and other fungal infections. The European Society of Clinical Microbiology and Infectious Disease (ESCMID) and the European Confederation of Medical Mycology (ECMM) joint guideline provides a comprehensive, evidence-based summary of antifungal therapy specifically for the treatment of mucormycosis. The 2016 IDSA guideline identifies voriconazole as the preferred primary therapy for most invasive syndromes of aspergillus. Other primary agents, depending on the syndrome: include liposomal amphotericin B, caspofungin, micafungin, and posaconazole. Alternative agents are also presented when primary agents cannot be used, depending on the syndrome present, include: liposomal amphotericin B, lipid complex amphotericin B, aerosolized amphotericin B, isavuconazole, caspofungin, micafungin, posaconazole, itraconazole, and voriconazole when it was not used as a primary agent. For mucormycosis the 2014 ESCMID/ECMM lists posaconazole or lipid formulations of amphotericin as first-line options for treatment of mucormycosis in certain types of individuals. The guideline lists posaconazole, lipid formulations of amphotericin B, and caspofungin for salvage treatment of mucormycosis in specific types of populations in adults. Posaconazole is recommended as prophylaxis for mucormycosis in neutropenia or graft-versus-host disease.

Isavuconazonium is a prodrug that is hydrolyzed in the blood by esterases, predominantly by butylcholinesterase, to the active component isavuconazole. Isavuconazole inhibits the synthesis of ergosterol, a key component of the fungal cell membrane, through the inhibition of cytochrome P-450 dependent enzyme lanosterol 14-alpha-demethylase. This enzyme is responsible for the conversion of lanosterol to ergosterol. An accumulation of methylated sterol precursors and a depletion of ergosterol within the fungal cell membrane weakens the membrane structure and function. Mammalian cell demethylation is less sensitive to isavuconazole inhibition.

---

### Resources:

Cresamba (isavuconazonium) product information, revised by Astellas Pharma US, Inc. 03-2020. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed May 06, 2022.



An Independent Licensee of the Blue Cross Blue Shield Association

**PHARMACY COVERAGE GUIDELINES**  
**SECTION: DRUGS**

**ORIGINAL EFFECTIVE DATE:** 5/19/2016  
**LAST REVIEW DATE:** 5/19/2022  
**LAST CRITERIA REVISION DATE:** 5/19/2022  
**ARCHIVE DATE:**

---

## **CRESEMBA® (isavuconazonium sulfate) oral**

---

Palmer SM, Zaas A, Messin JA. Fungal infections following lung transplantation. In: UpToDate, Blumberg EA, Trulock EP, Bond S (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Topic last updated November 29, 2021. Accessed May 06, 2022.

Wingard JR. Prophylaxis of invasive fungal infections in adults hematopoietic cell transplant recipients. In: UpToDate, Bow E, Bond S (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Topic last updated August 09, 2021. Accessed May 06, 2022.

Wingard JR. Prophylaxis of invasive fungal infections in adults with hematologic malignancies. In: UpToDate, Bow E, Bond S (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Topic last updated June 16, 2021. Accessed May 06, 2022.

Patterson TF. Treatment and prevention of invasive aspergillosis. In: UpToDate, Kauffman CA, Hall KK. (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Topic last update January 12, 2022. Accessed May 06, 2022.

Kauffman CA. Treatment of chronic pulmonary aspergillosis. In: UpToDate, Baddley JW, Hall KK (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Topic last updated November 17, 2020. Accessed May 06, 2022.

Cox GM. Mucormycosis (zygomycosis). In: UpToDate, Kauffman CA, Bond S (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Topic last update January 12, 2022. Accessed May 06, 2022.